



Sickle Cell Genetics Lab

Diagnosing Baby Marie™

Contents

Getting started

At a glance	P. 03
Class time requirements	P. 04
Materials needed	P. 05
Teacher prep	P. 07
Student workstation setup	P. 10

Student guide

Background information	P. 12
Today's lab	P. 16
Glossary	P. 21
Student lab protocol	P. 23
Post-lab questions	P. 24
CER table	P. 26
Extension: Sequence analysis of the sickle cell mutation	P. 28
Extension: Using the Hardy-Weinberg equation	P. 31
Extension: Sickle cell disease gene therapy	P. 33

Instructor guide

Expected results	P. 35
Unexpected results and troubleshooting	P. 36
Notes on lab design	P. 37
Additional student supports	P. 39
Extension activities	P. 39
Learning goals and skills developed	P. 40
Standards alignment	P. 40

At a glance

Lab overview

Initial testing for baby Marie suggests she might have sickle cell disease. Use gel electrophoresis to test baby Marie and her family for the sickle cell allele.

TECHNIQUES

Micropipetting
Gel electrophoresis

TOPICS

Genotype to phenotype
Inheritance
Biotechnology

LEVEL

General high school
Advanced high school
College

Planning your time


SINGLE CLASS: 45 MIN.

See the next page for detailed class time requirements


**Teacher prep
(30 min.)**

See page 7 for details. We recommend that the teacher prepare the gels and dispense the DNA samples before class.

**Gel
electrophoresis**



**Interpret
results**



Technical support

If you have any questions about implementing this activity, contact support@minipcr.com.

Class time requirements

This activity can be completed in a single 45-minute class period if the gels have been prepared in advance.

Steps	Time required
Prep Make gels	We recommend the teacher prepare the gels outside of class (see page 7). Allot 30 minutes of class time if you opt to have students prepare the gels.
1 Load gel	10 minutes
2 Run gel	15-20 minutes The gel does not need to be actively monitored during this time.
3 Interpret results	5 minutes

Materials needed

Supplied in kit (KT-1502-01)

- Kit contains DNA samples for eight lab groups.
- If kept in the freezer, reagents can be stored for 12 months after receipt. If kept in the refrigerator, reagents can be stored for 6 months after receipt.
- Reagents for preparing gels, plastic tubes for distributing samples to individual groups, and pipette tips are sold separately. See below for details.

Contents	Provided	Required per group	Storage
Simulated patient DNA samples <ul style="list-style-type: none"> • Jacqueline DNA • Kumar DNA • Lewis DNA • Marie DNA 	150 µl each	15 µl each	Freezer
Fast DNA Ladder 1	150 µl	15 µl	Freezer

Electrophoresis reagents and plastics sold separately

- This lab requires 2% agarose gels with a fluorescent DNA stain (e.g., SeeGreen™ or GelGreen®).
- The [Learning Lab Companion Kit](#) (KT-1510-01) provides sufficient reagents to prepare and run eight gels when using the blueGel or Bandit electrophoresis systems, as well as plastic tubes to distribute samples to student groups.
- Alternatively, [bulk electrophoresis reagents](#) and [plastics](#) (tubes, pipette tips) are available for purchase from miniPCR bio.
- Gel electrophoresis reagents and plastics can also be purchased from other suppliers.

Required equipment

- This lab is compatible with any horizontal gel electrophoresis system in combination with:
 - A fluorescent DNA stain (e.g., SeeGreen™ or GelGreen®).
 - A transilluminator that is compatible with the DNA stain used. Fluorescent DNA stains typically require blue light (~470 nm) or UV (~260 nm) illumination.
- The table below outlines gel electrophoresis equipment from miniPCR bio that meets these requirements:

Item	Recommended quantity
Gel electrophoresis and visualization system Option 1: blueGel™ OR GELATO™ electrophoresis systems with integrated blue light transilluminator Option 2: Bandit™ STEM electrophoresis kit paired with the Viewit™ Illumination Kit Option 3: Bandit™ STEM electrophoresis kit paired with a blueBox™ blue light transilluminator	1 blueGel can be shared by two groups 1 GELATO can be shared by four groups 1 Bandit + 1 Viewit per group 1 Bandit per group + 1 blueBox for the class to share
Micropipettes and tips 2-20 µl adjustable or 10 µl fixed volume	1 pipette per group

AVAILABLE AT MINIPCR.COM

Other materials supplied by user

- Distilled water
- Microwave or hot plate
- Heat-resistant flask or beaker
- Disposable laboratory gloves
- Protective eyewear
- Fine-tipped permanent marker

Teacher prep



Protective gloves and eyewear should be worn for the entirety of this experiment.

Overview

The table below provides an overview of the teacher prep, and the subsequent pages provide detailed instructions.

Prep	Time required	Timeline
Dispense reagents	10 minutes	Can be completed up to one week before use
Prepare electrophoresis buffer and agarose gels	20 minutes	Varies - If using gel reagents from miniPCR, gels can be prepared up to five days before use

Dispense reagents

- DNA samples can be dispensed up to one week in advance and stored in the refrigerator until use.
- This kit provides sufficient reagents for eight lab groups.

Materials needed

From the lab kit (stored in the freezer):

- Jacqueline DNA
- Kumar DNA
- Lewis DNA
- Marie DNA
- Fast DNA Ladder 1


Supplied by user:

- Plastic tubes for dispensing reagents (1.5 ml or 0.2 ml tubes can be used)
- 2-20 μ l micropipettes and tips
- Fine-tipped permanent marker

1. Thaw reagents by placing tubes at room temperature.
2. Collect the liquid at the bottom of each tube. Either spin briefly in a microcentrifuge or shake the liquid down with a flick of the wrist.
3. When you open each tube, check for liquid stuck inside the cap. If necessary, put the cap back on and repeat step 2.

4. For each lab group, dispense the following reagents into labeled plastic tubes. 1.5 ml or 0.2 ml plastic tubes can be used.

- | | |
|------------------------|------------|
| - Jacqueline DNA | 15 μ l |
| (label tube as "J") | |
| - Kumar DNA (tube K) | 15 μ l |
| - Lewis DNA (tube L) | 15 μ l |
| - Marie DNA (tube M) | 15 μ l |
| - DNA Ladder (tube DL) | 15 μ l |



5. If you are preparing the DNA samples more than 24 hours before class, store the tubes in the refrigerator until use. Dispensed DNA samples can be stored in the refrigerator for up to one week before use.

Prepare gel electrophoresis buffer and agarose gels

1. Prepare electrophoresis buffer.
 - Follow the manufacturer's instructions to prepare buffer solution.
 - The volume of buffer needed varies depending on the gel electrophoresis system.
 - For the blueGel and Bandit electrophoresis systems, 600 ml of TBE buffer is sufficient for at least eight gel runs.
 - For other systems, refer to the manufacturer's instructions for:
 - (1) The buffer volume needed to prepare agarose gels.
 - (2) The buffer volume needed for use as running buffer.
2. Prepare 2% agarose gels with fluorescent DNA stain.
 - Each group will need four lanes, plus one lane for ladder. If groups are sharing gels, a single lane for the ladder per gel is sufficient.
 - This lab kit is compatible with any molecular grade agarose and fluorescent DNA stain (e.g., SeeGreen™ or GelGreen®).
 - The volume of gel needed varies based on the gel electrophoresis system you are using. Refer to the manufacturer's instructions.
 - If using gel electrophoresis reagents from miniPCR bio, gels can be prepared up to five days in advance. Store prepared gels at room temperature in an airtight container protected from light. Do NOT soak the gels in buffer or wrap them in paper towels.

Detailed instructions for preparing buffer and gels for miniPCR electrophoresis systems



blueGel

<https://links.minipcr.com/gelpouring>



Bandit

<https://links.minipcr.com/BanditDNAgel>

Student workstation setup

At the start of this experiment, every lab group should have:

Simulated patient DNA samples <ul style="list-style-type: none"> • Jacqueline DNA (tube J) • Kumar DNA (tube K) • Lewis DNA (tube L) • Marie DNA (tube M) 	15 μ l each
Fast DNA Ladder 1 (tube DL)	15 μ l
2-20 μ l micropipette or 10 μ l fixed volume micropipette	1
Micropipette tips	At least 5
Electrophoresis buffer *Volume depends on your electrophoresis system	30 ml TBE if using a blueGel or Bandit
5 wells in a 2% agarose gel with fluorescent DNA stain	

Student guide



Background information	P. 12
Today's lab	P. 16
Glossary	P. 21
Student lab protocol	P. 23
Post-lab questions	P. 24
CER table	P. 26
Extension: Sequence analysis of the sickle cell mutation	P. 28
Extension: Using the Hardy-Weinberg equation	P. 31
Extension: Sickle cell disease gene therapy	P. 33




Background information

Sickle cell disease


1

- **Red blood cells** carry oxygen from the lungs to cells throughout our bodies.
- Normal red blood cells are disk-shaped and very flexible, allowing them to squeeze through tiny capillaries in our circulatory system.



2

- **Sickle cell disease** is a condition that causes red blood cells to become crescent-shaped and resemble a farm tool called a sickle.
- Sickled red blood cells clump together, creating blood flow blockages that can lead to infection, pain, or even death.






The sickle cell mutation


1

- **Hemoglobin** is a protein found in red blood cells that is responsible for transporting oxygen throughout the body.
- Hemoglobin is made of four protein subunits: two **alpha-globin (α -globin)** subunits and two **beta-globin (β -globin)** subunits.
- Hemoglobin proteins move freely inside the cytoplasm of red blood cells. This normal form of hemoglobin protein is called **hemoglobin A**.



2

- Sickle cell disease is caused by a **mutation** that substitutes a single **DNA nucleotide** in the *β-globin* gene.
- The mutation leads to a single amino acid substitution in the β -globin protein and causes a **hydrophobic** amino acid to face the watery cytoplasm.
- The mutated amino acids repel water and form hydrophobic links with each other, assembling long chains of hemoglobin proteins that distort the red blood cell into a sickle shape.
- This abnormal form of hemoglobin protein is called **hemoglobin S**.





Background: Stop and think


- Q1. What type of mutation causes sickle cell disease?
- Deletion of a DNA nucleotide.
 - Addition of a DNA nucleotide.
 - A single DNA nucleotide substitution.
 - A large chromosomal rearrangement.
- Q2. How does the sickle cell mutation affect the hemoglobin protein?
- It prevents hemoglobin from binding oxygen.
 - It prevents hemoglobin from being produced.
 - It causes hemoglobin proteins to cluster together.
 - It increases hemoglobin's ability to bind oxygen.



Inheritance of sickle cell disease

1

- There are two **alleles** of the β -globin gene relevant to sickle cell disease: the normal β -globin allele and the mutated sickle cell allele.
- It is standard to abbreviate the normal β -globin allele “A” (for hemoglobin A) and the sickle cell allele “S” (for hemoglobin S).
- As you can see in the pedigree chart below, there are three possible **genotypes** relevant to sickle cell disease.



2

- Only individuals with two copies of the sickle cell allele (SS) have sickle cell disease.
- This means that sickle cell disease follows a **recessive inheritance** pattern.

3

- Individuals with one copy of the normal β -globin allele and one copy of the sickle cell allele (AS) are said to have **sickle cell trait**.
- People with sickle cell trait have some hemoglobin clumping, but under normal conditions, their red blood cells do not sickle, and they do not experience symptoms.

4

- Individuals with two copies of the normal β -globin allele (AA) are unaffected.
- AA individuals can only pass down the normal β -globin allele.



Background: Stop and think

Q3. True or false: A person with sickle cell disease must carry two copies of the sickle cell allele.



Today's lab

Meet the Patel family

1

- You are a doctor working with the Patels, who recently welcomed baby Marie to their family.
- Marie's routine newborn screening revealed low levels of normal hemoglobin, suggesting sickle cell disease.



2

- **Goal: you will use DNA testing to determine whether Marie has sickle cell disease!**
- Because sickle cell disease is inherited, you will also test Marie's parents, Jacqueline and Kumar, and her older brother, Lewis, to determine if they carry the sickle cell allele.

Jacqueline: Jacqueline is a 32-year-old female. She is of primarily African descent, which can be a risk factor for carrying the sickle cell allele. Jacqueline suffers from occasional migraine headaches but is otherwise healthy.

Kumar: Kumar is a 32-year-old male born in India. Indian heritage can be a risk factor for carrying the sickle cell allele. Kumar reports nothing abnormal in his medical history.

Lewis: Lewis is a healthy 4-year-old boy and has had no major illnesses. Lewis did not show any abnormalities in his routine infant screening.

Marie: Marie is 2 months old, and her newborn screening suggests sickle cell disease.

Risk factors for carrying the sickle cell allele


If sickle cell disease is so dangerous, why is the sickle cell allele relatively common in some populations? It turns out that carrying the sickle cell allele confers protection against severe **malaria**, an infectious disease spread by mosquito bites. In regions where malaria is prevalent, there is **selective pressure** for the sickle cell allele to remain in the population. The sickle cell allele is more common in people with ancestry from eastern and central sub-Saharan Africa and certain areas of the Mediterranean, Middle East, and India. All of these regions have high rates of malaria.



DNA testing

1

- A DNA test can be used to diagnose sickle cell disease.
- DNA testing involves several steps. The following procedure will determine if baby Marie or her family members carry the sickle cell allele.



2

A blood or tissue sample is taken from the patient. DNA is then collected from that sample.

3

Then, a method called **PCR** makes billions of copies of a section of the β -globin gene.

4

Next, a step to identify the sickle cell mutation is performed. This is explained on the next page.

5

You will perform the last step of the DNA test to determine if the mutated sickle cell allele is present.

Living with sickle cell disease

What if baby Marie tests positive for sickle cell disease? Advances in modern medicine allow most patients to manage sickle cell disease. Previously, standard treatments only addressed symptoms, with bone marrow transplants as the sole cure. This procedure replaces a patient's blood stem cells with healthy donor cells, producing normal red blood cells. However, finding a compatible donor is challenging. Because sickle cell disease is a genetic disease, it can be treated by modifying a patient's DNA. Recent advances have led to the first approved gene therapy treatments for patients with sickle cell disease. For more information, refer to https://links.minipcr.com/crispr_sicklecell.



Identifying the sickle cell allele using a restriction enzyme

1

- In this lab, PCR is used to copy a 400 bp segment of the *β-globin* gene that includes the location of the sickle cell mutation.
- Then, a **restriction digestion** is used to identify if the sickle cell mutation is present.

2


- **Restriction enzymes** recognize and cut specific short **DNA sequences**. This lab uses a restriction enzyme that cuts a DNA sequence found only in the normal *β-globin* allele.
- The sequence CTGAG in the normal *β-globin* allele is cut by the restriction enzyme **Ddel**.

3

- The sickle cell mutation changes an adenine (A) to a thymine (T).
- This change means the enzyme Ddel does not cut the sickle cell allele.
- The difference in whether the DNA was cut or not can be observed using gel electrophoresis, which is explained on the next page.


Normal *β-globin* allele

Ddel cuts DNA



Sickle cell allele

Ddel does NOT cut DNA



Background: Stop and think

Q4. What is the role of PCR in this experiment?

- Copy a section of the normal *β-globin* allele.
- Copy a section of the sickle cell allele.
- Copy a section of the normal *β-globin* allele and the sickle cell allele.
- None of the above.

Q5. What is the role of the restriction digestion in this experiment?


- Confirm successful DNA extraction.
- Read the sequence of the *β-globin* gene.
- Confirm successful PCR.
- Differentiate between the normal *β-globin* allele and the sickle cell allele.



Interpreting gel electrophoresis results

1

- Recall that PCR was used to copy a 400 bp segment of the β -globin gene and then restriction digestion was performed.
- The sickle cell allele is not cut by the restriction enzyme, while the normal β -globin allele is cut.
- You will use gel electrophoresis to view the results and determine if the Patels carry the sickle cell allele.
- Gel electrophoresis** separates pieces of DNA by size, with smaller pieces of DNA traveling farther.



2

Individuals who have sickle cell disease (SS) will show a single 400 bp band on the gel. This band corresponds to the sickle cell allele, which is not cut by the restriction enzyme.

3

Unaffected individuals (AA) will show two bands. These bands correspond to the normal β -globin allele, which is cut by the restriction enzyme into two fragments: 250 bp and 150 bp.

4

Individuals with sickle cell trait will show three bands: the 400 bp band that corresponds to the sickle cell allele, as well as the 250 bp and 150 bp bands that correspond to the normal β -globin allele.




Background: Stop and think

- Q6. Why is gel electrophoresis a good tool for interpreting the results of a restriction digest?
- A. Because it allows you to make copies of DNA.
 - B. Because it allows you to extract DNA from cells.
 - C. Because it allows you to separate DNA fragments by size.
 - D. Because it allows you to cut specific DNA sequences.

Imagine getting the gel electrophoresis result shown below after testing a group of patients for sickle cell disease.

- Q7. Based on the gel results, which individual(s) would you diagnose with sickle cell trait?
- A. Person A
 - B. Person B
 - C. Person C
 - D. Person D
 - E. Person E
 - F. Person F



- Q8. Based on the gel results, which individual(s) would you diagnose with sickle cell disease?
- A. Person A
 - B. Person B
 - C. Person C
 - D. Person D
 - E. Person E
 - F. Person F



Glossary

Red blood cell: A type of blood cell that delivers oxygen to cells throughout the body. Normal red blood cells are disk-shaped and quite flexible, allowing them to squeeze through small capillaries.

Sickle cell disease: A genetic disease where red blood cells take on a sickle shape. Sickled red blood cells cause blood flow blockages, leading to episodes of severe pain, organ failure, or even death.

Hemoglobin: An oxygen-binding protein found in red blood cells. Hemoglobin is composed of two α -globin subunits and two β -globin subunits.

Alpha-globin (α -globin): One of the protein subunits in the hemoglobin protein.

Beta-globin (β -globin): One of the protein subunits in the hemoglobin protein. A mutation in the *β -globin* gene causes sickle cell disease.

Hemoglobin A: Normal hemoglobin protein.

Mutation: A change or variation in the DNA sequence. Mutations can have beneficial, neutral, or harmful effects.

DNA nucleotide: A building block of DNA. Each nucleotide consists of a phosphate group, a sugar, and a nitrogenous base. There are four DNA bases: adenine (A), cytosine (C), guanine (G), and thymine (T).

Hydrophobic: Lacking an affinity for water. Hydrophobic amino acids are repelled by the watery cytoplasm of the cell.

Hemoglobin S: Abnormal hemoglobin protein in individuals with sickle cell disease. Molecules of hemoglobin S clump together and can lead to the formation of long strands that distort the shape of the red blood cell into a sickle.

Allele: One of two or more alternative versions of the same gene. Different alleles of the same gene have differences in the DNA sequence.

Genotype: An organism's genetic makeup. For a specific gene, an organism's genotype is usually a pair of alleles, one inherited from each biological parent.

Recessive inheritance: A pattern of inheritance where a trait is only present when the organism carries two copies of the same allele.



Sickle cell trait: A condition in which individuals carry one copy of the sickle cell allele. People with the sickle cell trait produce some abnormal hemoglobin protein but usually do not exhibit symptoms of sickle cell disease. However, their red blood cells can sickle under extreme environmental conditions, including low oxygen levels.

Malaria: An infectious disease caused by a parasite carried by certain species of mosquitoes. Malaria is common in certain regions where these mosquitoes are prevalent. Severe cases of malaria can lead to death. Carrying the sickle cell allele provides some protection against severe malaria infections.

Selective pressure: Environmental factors that influence an organism's ability to survive and reproduce, favoring individuals with certain traits. In areas where malaria is common, there is selective pressure for the sickle cell allele to remain in the population, even though carrying two copies of the sickle cell allele is disadvantageous to the individuals affected by sickle cell disease.

Polymerase Chain Reaction (PCR): A technique used to make multiple copies of a specific DNA segment for further study. For more detailed information on PCR, refer to <https://www.minipcr.com/polymerase-chain-reaction/>.

Restriction digestion: The use of a restriction enzyme to cut, or digest, a DNA sample.

Restriction enzyme: An enzyme that recognizes a specific, short DNA sequence (typically 4-8 base pairs long) and cuts the DNA at that location.

DNA sequence: The order of nucleotides, or bases, in a DNA molecule.

Ddel: A restriction enzyme that specifically recognizes and cuts a short DNA sequence present in the normal *β-globin* allele and absent in the sickle cell allele.

Gel electrophoresis: A method that separates pieces of DNA by length. For more detailed information on gel electrophoresis, refer to <https://www.minipcr.com/gel-electrophoresis/>.



Student lab protocol

! Protective gloves and eyewear should be worn for the entirety of this experiment.

1. Place the prepared gel into the electrophoresis chamber.
2. Add enough electrophoresis buffer to fill the chamber and just cover the gel.
 - You will need 30 ml of TBE buffer for a blueGel or Bandit electrophoresis system. Do not overfill the chamber.
 - If using another electrophoresis system, refer to the manufacturer's instructions for the recommended buffer type and volume.
3. Use a micropipette to load samples in the following order. To prevent contamination, use a new tip for each sample.
 - Well 1: 10 μ l DNA Ladder (tube DL)
 - Well 2: 10 μ l Jacqueline DNA (tube J)
 - Well 3: 10 μ l Kumar DNA (tube K)
 - Well 4: 10 μ l Lewis DNA (tube L)
 - Well 5: 10 μ l Marie DNA (tube M)
4. Run the gel for 15-20 minutes.
 - The blueGel and Bandit electrophoresis systems run at a fixed voltage.
 - If using another gel electrophoresis system, set the voltage in the 70-90 V range.
5. To visualize the DNA samples, turn on the blue light in your electrophoresis system, or move the gel to a transilluminator.
6. If needed, continue to run the gel until there is sufficient separation between the 100-500 bp bands in the ladder to interpret the results.
7. If desired, take a photo to document the results.
8. Compare the bands from the DNA samples to the DNA ladder to obtain size estimates.

Detailed operating instructions for miniPCR electrophoresis systems



blueGel

<https://links.minipcr.com/blueGelRun>



Bandit

<https://links.minipcr.com/BanditViewit>





Post-lab questions

Interpreting results

1. Use the schematic gel on the right to draw what your results look like. For each sample, draw the bands that you see on your actual gel.
2. Label the bands with an approximate size (in base pairs). Use the image of the ladder on the previous page to help you.
3. Use your gel electrophoresis results to complete the table below.



- A. Use checkmarks to record the gel electrophoresis results in the first two rows of the table.
- B. Record each person's genotype and diagnosis.

	Jacqueline	Kumar	Lewis	Marie
Sickle cell allele (400 bp)				
Normal <i>β-globin</i> allele (150 + 100 bp)				
Genotype (AA, AS, SS)				
Diagnosis (Unaffected, sickle cell trait, sickle cell disease)				




Critical thinking

4. Imagine Jacqueline and Kumar have a third child. Fill out the Punnett square to help you answer the following questions.

- What is the chance this child will have two copies of the normal β -globin gene?
- What is the chance this child will have sickle cell trait?
- What is the chance this child will have sickle cell disease?


Cross: _____ x _____
 Jacqueline's genotype Kumar's genotype

Jacqueline



5. Fill in the pedigree chart for baby Marie's family. Kumar's parents have been filled in for you as an example.

- Record each family member's genotype.
- Shade each family member's shape if appropriate to indicate the presence of the sickle cell allele.





CER table

Fill in the table based on your results from the lab. Refer to the rubric on the next page.

Question:

Does baby Marie have sickle cell disease?

Claim

Make a clear statement that answers the above question.

Evidence

Provide data from the lab that supports your claim.

Reasoning

Explain clearly why the data you presented supports your claim. Include the underlying scientific principles that link your evidence to your claim.



Score	4	3	2	1
CLAIM A statement that answers the original question/problem.	Makes a clear, accurate, and complete claim.	Makes an accurate and complete claim.	Makes an accurate but incomplete or vague claim.	Makes a claim that is inaccurate.
EVIDENCE Data from the experiment that supports the claim. Data must be <u>relevant</u> and <u>sufficient</u> to support the claim.	All of the evidence presented is highly relevant and clearly sufficient to support the claim.	Provides evidence that is relevant and sufficient to support the claim.	Provides relevant but insufficient evidence to support the claim. May include some non-relevant evidence.	Only provides evidence that does not support claim.
REASONING Explain why your evidence supports your claim. This must include scientific principles/knowledge that you have about the topic to show why the data counts as evidence.	Provides reasoning that clearly links the evidence to the claim. Relevant scientific principles are well integrated in the reasoning.	Provides reasoning that links the evidence to the claim. Relevant scientific principles are discussed.	Provides reasoning that links the evidence to the claim, but does not include relevant scientific principles or uses them incorrectly.	Provides reasoning that does not link the evidence to the claim. Does not include relevant scientific principles or uses them incorrectly.

We recommend that teachers use the following scale when assessing this assignment using the rubric. Teachers should feel free to adjust this scale to their expectations.

Rubric score	3	4	5	6	7	8	9	10	11	12
Equivalent	55	60	65	70	75	80	85	90	95	100



Extension: Sequence analysis of the sickle cell mutation

Restriction enzyme analysis

Recall that this activity uses the restriction enzyme DdeI to differentiate between the normal β -globin allele and the sickle cell allele. DdeI recognizes the following sequence:



The “N” means that the middle nucleotide of the sequence could be any of the four DNA nucleotides: A, T, C, or G. The sequences CTAAG, CTTAG, CTGAG, and CTCAG would all be cut by DdeI. The red line indicates exactly where the cut happens in the DNA strands.

The sickle cell mutation is a single nucleotide substitution near the start of the β -globin gene. Shown below are the first 30 base pairs of the normal β -globin allele and the sickle cell allele.

	Coding sequence nucleotide 1	Coding sequence nucleotide 10	Coding sequence nucleotide 20	Coding sequence nucleotide 30
	↓	↓	↓	↓
Seq. 1	$ \begin{array}{l} 5' \text{ A T G G T G C A T C T G A C T C C T G A G G A G A A G T C T } 3' \\ 3' \text{ T A C C A C G T A G A C T G A G G A C T C C T C T T C A G A } 5' \end{array} $			
Seq. 2	$ \begin{array}{l} 5' \text{ A T G G T G C A T C T G A C T C C T G T G G A G A A G T C T } 3' \\ 3' \text{ T A C C A C G T A G A C T G A G G A C A C C T C T T C A G A } 5' \end{array} $			

1. Circle the single base pair that differs between Sequence 1 and Sequence 2.
2. Find the DdeI recognition sequence and draw a box around it.
3. DdeI only cuts the normal β -globin allele. Based on where you identified the cut site for DdeI, is sequence 1 or sequence 2 the normal β -globin allele?



Mutation analysis

The sequences you analyzed on the previous page code for the first 10 amino acids of the β -globin protein. Now you will transcribe and translate the sequence to examine the effect of the sickle cell mutation on the β -globin protein sequence.

- Transcribe each DNA sequence into mRNA.
 - To help you, the first 30 nucleotides from the template strand have been broken up into codons.
 - As an example, the first codon in Sequence 1 has been filled in for you.
- Use the mRNA codon table on the next page to translate the mRNA into protein. As an example, the first amino acid in Sequence 1 has been filled in for you.

Seq. 1

		Coding sequence nucleotide 1		Coding sequence nucleotide 10		Coding sequence nucleotide 20		Coding sequence nucleotide 30			
		↓		↓		↓		↓			
DNA	3'	TAC	CAC	GTA	GAC	TGA	GGA	CTC	CTC	TTC	AGA
mRNA	5'	AUG	---	---	---	---	---	---	---	---	---
Amino acid		Met	---	---	---	---	---	---	---	---	---

Seq. 2

		Coding sequence nucleotide 1		Coding sequence nucleotide 10		Coding sequence nucleotide 20		Coding sequence nucleotide 30			
		↓		↓		↓		↓			
DNA	3'	TAC	CAC	GTA	GAC	TGA	GGA	CAC	CTC	TTC	AGA
mRNA	5'	---	---	---	---	---	---	---	---	---	---
Amino acid		---	---	---	---	---	---	---	---	---	---

- Describe the change that occurred in the amino acid sequence.
 - Circle the amino acid position in the diagrams above affected by the sickle cell mutation.
 - What amino acid does the normal β -globin allele have in that position?
 - What amino acid does the sickle cell allele have in that position?



- Summarize why this single amino acid substitution leads to the mutated β -globin protein sticking together. Refer to the background information on page 13 if you need a reminder of the properties of the amino acids involved.

mRNA codon table

		Second Position Nucleotide									
		U		C		A		G			
First Position Nucleotide	U	UUU	Phenylalanine (Phe, F)	UCU	Serine (Ser, S)	UAU	Tyrosine (Tyr, Y)	UGU	Cysteine (Cys, C)	U	
		UUC		UCC		UAC		UGC		C	
		UUA	Leucine (Leu, L)	UCA		STOP	UAA	UGA	STOP	A	
		UUG		UCG			UAG			UGG	Tryptophan (Trp, W)
	C	CUU	Leucine (Leu, L)	CCU	Proline (Pro, P)		CAU	Histidine (His, H)	CGU	Arginine (Arg, R)	U
		CUC		CCC			CAC		CGC		C
		CUA		CCA		CAA	Glutamine (Gln, Q)	CGA	A		
		CUG		CCG		CAG		CGG	G		
	A	AUU	Isoleucine (Ile, I)	ACU	Threonine (Thr, T)	AAU	Asparagine (Asn, N)	AGU	Serine (Ser, S)	U	
		AUC		ACC		AAC		AGC		C	
		AUA		ACA		AAA	Lysine (Lys, K)	AGA	Arginine (Arg, R)	A	
		AUG	Methionine (Met, M) START	ACG		AAG		AGG		G	
	G	GUU	Valine (Val, V)	GCU	Alanine (Ala, A)	GAU	Aspartic Acid (Asp, D)	GGU	Glycine (Gly, G)	U	
		GUC		GCC		GAC		GGC		C	
		GUA		GCA		GAA	Glutamic Acid (Glu, E)	GGA		A	
		GUG		GCG		GAG		GGG		G	
		Third Position Nucleotide									



Extension: Using the Hardy-Weinberg equation

In a population that is not evolving, the prevalence of different alleles within a population are expected to remain the same over time. A population in this state is referred to as being in Hardy-Weinberg equilibrium.

The Hardy-Weinberg equation describes the genotype frequencies you would expect to see in a population that is in Hardy-Weinberg equilibrium. In its simplest form, the Hardy-Weinberg equation describes a gene with two alleles. The convention is to use the symbols p and q to represent the frequency of each allele.

For a gene with two alleles, $p + q = 1$

For examining sickle cell disease:

p = Normal β -globin allele

q = Sickle cell allele

The Hardy-Weinberg equation predicts the genotypic frequencies for a population that is not evolving:

$$p^2 + 2pq + q^2 = 1$$

p^2 = the genotype frequency for individuals homozygous for the p allele

$2pq$ = the genotype frequency for heterozygous individuals

q^2 = the genotype frequency for individuals homozygous for the q allele

1. The frequency of the sickle cell allele in African American populations is thought to be close to 0.04. Use this information to answer the following questions. You can use a calculator but show your work.
 - A. What is the frequency of the normal β -globin allele?
 - B. What is the predicted genotype frequency for individuals who have two copies of the normal β -globin allele?
 - C. What is the predicted genotype frequency for individuals with sickle cell trait?
 - D. What is the predicted genotype frequency for individuals with sickle cell disease?



2. In Nigeria, it is estimated that 3% of all newborn babies have sickle cell disease. Use this information to estimate the percentage of the population you expect to have sickle cell trait. You can use a calculator but show your work.

3. Remember that the Hardy-Weinberg equation assumes that a population is in Hardy-Weinberg equilibrium and makes five assumptions about the population:

- The population size is infinite (or very large).
- There is no net migration into or out of the population.
- There is no mutation at the locus being tested.
- Mating in the population is random.
- There is no natural selection on the alleles being tested.

Based on these criteria, would you expect the normal β -globin allele and the sickle cell allele to be in perfect Hardy-Weinberg equilibrium in the global population? Explain your reasoning.



Extension activity: Sickle cell disease gene therapy

Use a paper-based modeling activity to explore the first use of CRISPR/Cas genome editing to treat sickle cell disease in human patients. Visit https://links.minipcr.com/crispr_sicklecell.






Instructor guide



Expected results	P. 35
Unexpected results and troubleshooting	P. 36
Notes on lab design	P. 37
Additional student supports	P. 39
Extension activities	P. 39
Learning goals and skills developed	P. 40
Standards alignment	P. 40



Expected results



This image represents results obtained after a 20 minute run using a blueGel electrophoresis system.

	Jacqueline	Kumar	Lewis	Marie
Sickle cell allele (400 bp)	✓	✓		✓
Normal β -globin allele (250 + 150 bp)	✓	✓	✓	
Genotype	AS	AS	AA	SS
Diagnosis	Sickle cell trait	Sickle cell trait	Unaffected	Sickle cell disease

For technical support, contact support@minipcr.com

For answers to the student questions, email answers@minipcr.com

Please include in the body of the email:

- The name of the lab
- Your name, school, and job title



Unexpected results and troubleshooting

If **fluorescent DNA bands are not visible on the gel**, the following may have occurred:

- Failure to use a fluorescent DNA stain: This lab requires agarose gels made with a fluorescent DNA stain (e.g., SeeGreen™ or GelGreen®). DNA stains that reveal DNA with a visible blue compound are less sensitive and are not compatible with this lab kit.
- Incorrect visualization conditions: Fluorescent DNA stains (e.g., SeeGreen™ or GelGreen®) must be viewed using a blue light or UV transilluminator. The blueGel system has an integrated blue light transilluminator. For DNA visualization, ensure that you have turned on the blueGel's blue light by pressing the light bulb button.
- Samples were run off the gel: If you run the gel too long, DNA samples may migrate off the gel. Monitor progress by occasionally checking the DNA samples under a transilluminator or tracking the loading dye, which is visible to the eye. Stop the run before the colored loading dye reaches the end of the gel.
- Reagents were stored improperly and/or are expired: DNA samples can be stored in the freezer for up to twelve months after receipt or in a refrigerator for six months. Storage under different conditions or in excess of this guidance may impair performance.

If **some or all of the bands appear faint**, the following may have occurred:

- Failure to load the DNA samples: Loading DNA samples for gel electrophoresis takes a little practice. The bands will appear faint if students do not successfully deposit the full sample volume into the well. Refer to <https://www.minipcr.com/how-to-load-a-gel-electrophoresis/> for gel loading tips.
- Non-optimal visualization conditions: Dimming the lights in the room can make the fluorescent DNA stain easier to see in the transilluminator. If using the blueGel, viewing the gel using the Fold-a-View documentation hood and a smartphone camera will provide the best results.
- Old or improperly stored gels: Agarose gels can generally be prepared in advance, but the storage time and conditions depend on the fluorescent DNA stain being used. If using SeeGreen™ or GelGreen® DNA stain, gels can be prepared up to five days in advance. Store gels at room temperature in an airtight container protected from light. Do not soak the gels in buffer or wrap them in paper towels.
- Old or incorrectly prepared buffer: Gel visualization defects not directly ascribable to other causes may be remedied by using freshly prepared buffer compatible with your gel electrophoresis system. If using the blueGel, TBE buffer is recommended.

For tips on picture-perfect gels, see <https://www.minipcr.com/gel-electrophoresis-troubleshooting/>.

For additional technical support, contact support@minipcr.com.



Notes on lab design

This lab serves as an introduction to the relationship between genotype and phenotype. We believe this approach provides the right balance between intellectual engagement, inquiry, and accessibility. The design of this lab has simplified certain elements to achieve these goals. Some of these elements include:

- This lab uses prepared DNA to simulate the results of PCR amplification and restriction digest of a portion of the *β-globin* gene from human genomic DNA.
- While restriction digestion analysis is a powerful tool that is used in genetics in the way described in this lab, today, the relatively low cost of sequencing means that genetic testing for the sickle cell allele typically relies on DNA sequencing.
- While “dominant” and “recessive” are often used in biology textbooks and presented as a very straightforward concept, the relationship between alleles of a gene and the phenotypes they contribute to is often complicated. The sickle cell allele is a good example of this. On the next page, we explain how the inheritance pattern of the sickle cell allele changes depending on the trait being examined.



Sickle cell allele inheritance patterns

Sickle cell disease: In this activity, we focused on how the sickle cell allele causes sickle cell disease. Only people with two copies of the sickle cell allele (SS genotype) have the sickle cell disease phenotype. Therefore, sickle cell disease follows a recessive inheritance pattern.

Sickle cell disease: recessive inheritance pattern			
Genotype	AA	AS	SS
Phenotype	NO sickle cell disease	NO sickle cell disease	Sickle cell disease

But, the sickle cell allele also determines two other phenotypes that follow different inheritance patterns!

Hemoglobin type: The normal *β-globin* allele leads to the production of normal hemoglobin A, and the sickle cell allele leads to the production of hemoglobin S, which clumps together. We all have two alleles for the *β-globin* gene, one inherited from each parent, and both alleles are used to produce protein. That means that heterozygous individuals (genotype AS) have some normal hemoglobin A and some abnormal hemoglobin S. Because the effects of both alleles are observable, hemoglobin type follows a co-dominant inheritance pattern.

Hemoglobin type: co-dominant inheritance pattern			
Genotype	AA	AS	SS
Phenotype	Hemoglobin A protein	Hemoglobin A protein & Hemoglobin S protein	Hemoglobin S protein

Malaria resistance: People who carry the sickle cell allele produce hemoglobin S, which makes it harder for malaria parasites to reproduce within red blood cells. The presence of hemoglobin S provides protection from severe malaria infection, and having a single copy of the sickle cell allele confers this protection. This means that malaria resistance follows a dominant inheritance pattern.

Hemoglobin type: co-dominant inheritance pattern			
Genotype	AA	AS	SS
Phenotype	NO malaria resistance	Malaria resistance	Malaria resistance



Additional student supports

E-worksheets: The student questions accompanying this lab are available for download [here](#) as editable text documents you can customize and upload to your LMS. E-worksheets can also be accessed from the Curriculum Downloads tab at <https://www.minipcr.com/product/sickle-cell-genetics-lab/>.

miniPCR tutorials: Access an extensive set of free resources to help your students succeed in molecular biology techniques. Visit <https://www.minipcr.com/tutorials/>. The resources most relevant to this lab are listed below.

- **Micropipetting:** Video, worksheet, and hands-on activity resources to train students in the basic use of a micropipette.
- **Gel electrophoresis:** Video and worksheet activity instructing students on the fundamentals and practice of agarose gel electrophoresis.
- **PCR:** While students do not perform PCR in this lab, the samples they analyze represent PCR products. If you want to discuss PCR in more detail with your students, we have a video and worksheet activity instructing students on the fundamentals and practice of PCR.

miniPCR Digital: Interactive tools for experiment-based learning with or without hands-on lab kits. Visit <https://digital.minipcr.com/>.

Extension activities

The following optional extension activities are provided for students to explore topics more deeply.

Sequence analysis of the sickle cell mutation (page 28): Examine how the single nucleotide change in the *β-globin* gene affects hemoglobin's function and causes sickle cell disease symptoms.

Using the Hardy-Weinberg equation (page 31): Practice Hardy-Weinberg calculations using real-world data on *β-globin* allele frequencies.

Sickle cell disease gene therapy: Use a paper-based activity to model the first use of CRISPR/Cas genome editing to treat sickle cell disease in human patients. This activity contains all the background information on CRISPR/Cas genome editing for students to understand this cutting-edge treatment for sickle cell disease. https://links.minipcr.com/crispr_sicklecell.



Learning goals and skills developed

Student learning goals

- Correlate genotype and phenotype
- Predict genotype and phenotype using Punnett squares
- Apply basic probability rules to genetic analysis
- Identify how amino acid composition affects protein structure
- Relate changes in DNA and amino acid sequence to human disease
- Solve real-world problems using genetic analysis

Scientific inquiry skills

- Identify or pose a testable question
- Follow detailed experimental protocols
- Interpret data presented in a chart or table
- Make a claim based in scientific evidence
- Use reasoning to justify a scientific claim

Molecular biology skills

- Micropipetting
- Agarose gel electrophoresis

Standards alignment

The standards alignment document for this activity is available for download [here](#). This document can also be accessed from the Curriculum Downloads tab at <https://www.minipcr.com/product/sickle-cell-genetics-lab/>.

This activity is aligned to the following standards:

- Next Generation Science Standards: High School Life Science
- Advanced Placement Biology
- Texas Essential Knowledge and Skills: Biology
- Texas Essential Knowledge and Skills: Biotechnology
- Biotechnician Assistant Credentialing Exam
- Common Core ELA/Literacy Standards (9-10)

For additional information on alignment to state standards, please contact support@minipcr.com.